

Regulation and Function of Heat-Inducible Genes in *Bacillus subtilis*

WOLFGANG SCHUMANN, MICHAEL HECKER, AND TAREK MSADEK

25

In their natural environment, bacteria are frequently exposed to large and rapid temperature changes and, therefore, bacterial cells are equipped with a genetically regulated program to adapt to these temperature changes ensuring growth and survival. The best studied of these programs, the heat shock response, deals with the adaptation to a sudden increase in temperature, which is an important homeostatic mechanism that enables cells from animals, plants, and bacteria to survive a variety of environmental stresses besides heat stress (48, 49). It is characterized by the transiently increased synthesis of a number of proteins collectively designated heat shock proteins (hsps). The strong evolutionary conservation of the heat shock response argues that this response is beneficial for many kinds of cells and has evolved in order to detect and deal with the presence of unfolded, misfolded, damaged, or aggregated polypeptide chains. Furthermore, while some hsps have essential roles in the synthesis, transport, and folding of proteins and are often referred to as molecular chaperones (23), others are ATP-dependent proteases sometimes acting in concert with chaperones (24). In prokaryotes, the major hsps are encoded by single genes expressed at a basal level at all temperatures. Following a temperature upshift, the rates of expression of these genes abruptly accelerate and in general gradually decrease after 5 to 10 min. All bacterial heat shock genes are regulated at the level of transcription either by alternate sigma factors or by repressor proteins (for reviews, see references 27 and 92).

Two major groups of hsps constitute the heat stress stimulon of *Bacillus subtilis*: σ^B -dependent general stress proteins (also known as class II hsps and described in chapter 26), which are induced not only by heat but by a different set of stress and starvation stimuli conferring a nonspecific multiple stress resistance, and heat-specific stress proteins, which may exert a specific protective function against heat stress only. HrcA- and CtsR-dependent proteins (known as class I and class III hsps, respectively) belong to this heat-specific group, but ClpP and ClpC may have an intermediate role between heat-specific and general stress protein because of their double stress control.

CLASS I HEAT SHOCK GENES: THE HrcA REGULON

Discovery

Using gene probes, the *dnaK* and *groEL* genes were first identified and then cloned and sequenced (42, 68, 77, 87). The *groE* operon turned out to be bicistronic, consisting of *groES* and *groEL* (Fig. 1A). This genomic organization is highly conserved among eubacteria and has been described for all species studied so far; some may contain either an additional copy of *groEL*, such as *Streptomyces* spp. (45), and others may inherit up to five copies of the complete *groESL* operon, as described for *Bradyrhizobium japonicum* (18).

Sequencing of the region adjacent to *dnaK* revealed that it is flanked by *grpE* and *dnaJ*, and upstream of *grpE*, a fourth gene named *orf39* was identified coding for a protein of unknown function which turned out to encode the repressor of both the *dnaK* and *groESL* operons (see below). Sequencing of the whole region disclosed three additional *orfs* termed *orf35*, *orf28*, and *orf50* (29). Therefore, the complete *dnaK* operon is heptacistronic (Fig. 1B). While the function of *orf28* and *orf50* remains elusive, the deduced protein sequence of *orf35* exhibits significant homology to the ribosomal protein L11 methyltransferase encoded by *prmA* of *Escherichia coli* (81). The genomic organization of the *dnaK* operons of *Bacillus stearothermophilus*, *Clostridium acetobutylicum*, and *Staphylococcus aureus* is comparable to that of *B. subtilis* (5, 28, 56, 60).

The HrcA-CIRCE Regulation Mechanism

Both the *dnaK* and *groE* operons are preceded by a σ^A -type promoter that is used before and after heat stress (42, 68, 77, 87). Since σ^A is the major sigma factor produced constitutively, some other mechanism(s) must mediate the heat-inducible expression of these two operons. The first hint toward the elucidation of the regulation mechanism was the observation of a perfect inverted repeat of a length of 9 bp separated by a 9-bp spacer, located in both operons between the transcriptional and translational start sites. Since this

DegS/DegU two-component system, there is as yet no evidence to suggest that these regulators play a role in the heat shock induction of *sacB* expression.

The *ykoZ* gene, encoding a putative sigma factor, is strongly induced by heat stress in rich medium, but surprisingly is not induced in synthetic medium. Other stress and starvation signals tested so far do not induce *ykoZ*. It is interesting to note that *ykoZ* mutants grow poorly at 56°C on agar plates, but the *ykoZ*-dependent heat shock proteins that might be responsible for this growth defect have not yet been discovered (93).

T.M. is grateful to Georges Rapoport, in whose laboratory part of this work was carried out, and to The Institute for Genomic Research, the University of Oklahoma Advanced Center for Genome Technology, the Sanger Centre, and Genome Therapeutics for generously providing access to unfinished microbial genome sequences. W.S. and M.H. thank the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie for continuous financial support.

REFERENCES

- Ang, D., K. Liberek, D. Skowrya, M. Zylicz, and C. Georgopoulos. 1991. Biological role and regulation of the universally conserved heat shock proteins. *J. Biol. Chem.* 266:24233–24236.
- Antelmann, H., S. Engelmann, R. Schmid, and M. Hecker. 1996. General and oxidative stress responses in *Bacillus subtilis*: cloning, expression, and mutation of the alkyl hydroperoxide reductase operon. *J. Bacteriol.* 178:6571–6578.
- Bardwell, J. C. A., and E. A. Craig. 1987. Eukaryotic Mr 83,000 heat shock protein has a homologue in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 84:5177–5181.
- Bardwell, J. C. A., and E. A. Craig. 1988. Ancient heat shock gene is dispensable. *J. Bacteriol.* 170:2977–2983.
- Behrens, S., F. Narberhaus, and H. Bahl. 1993. Cloning, nucleotide sequence and structural analysis of the *Clostridium acetobutylicum* *dnaJ* gene. *FEMS Microbiol. Lett.* 114:53–60.
- Bernhardt, J., U. Völker, A. Völker, H. Antelmann, R. Schmid, H. Mach, and M. Hecker. 1997. Specific and general stress proteins in *Bacillus subtilis*—a two-dimensional protein electrophoresis study. *Microbiology* 143:999–1017.
- Bukau, B., and G. C. Walker. 1989. Cellular defects caused by deletion of the *Escherichia coli* *dnaK* gene indicate roles for heat shock protein in normal metabolism. *J. Bacteriol.* 171:2337–2346.
- Chastanet, A. Unpublished data.
- Chastanet, A., and T. Msadek. Unpublished data.
- Derré, I. Unpublished data.
- Derré, I., G. Rapoport, K. Devine, M. Rose, and T. Msadek. 1999. ClpE, a novel type of HSP100 ATPase, is part of the CtsR heat shock regulon of *Bacillus subtilis*. *Mol. Microbiol.* 32:581–593.
- Derré, I., G. Rapoport, and T. Msadek. 1999. CtsR, a novel regulator of stress and heat shock response, controls *clp* and molecular chaperone gene expression in Gram-positive bacteria. *Mol. Microbiol.* 31:117–131.
- Derré, I., G. Rapoport, and T. Msadek. Unpublished data.
- Deuerling, E., A. Mogk, C. Richter, M. Purucker, and W. Schumann. 1997. The *ftsH* gene of *Bacillus subtilis* is involved in major cellular processes such as sporulation, stress adaptation and secretion. *Mol. Microbiol.* 23:921–933.
- Deuerling, E., B. Paeslack, and W. Schumann. 1995. The *ftsH* gene of *Bacillus subtilis* is transiently induced after osmotic and temperature upshock. *J. Bacteriol.* 177:4105–4112.
- Duchêne, A.-M., C. J. Thompson, and P. Mazodier. 1994. Transcriptional analysis of *groEL* genes in *Streptomyces coelicolor* A3(2). *Mol. Gen. Genet.* 245:61–68.
- Fabret, C., and J. A. Hoch. 1998. A two-component signal transduction system essential for growth of *Bacillus subtilis*: implications for anti-infective therapy. *J. Bacteriol.* 180:6375–6383.
- Fischer, H. M., M. Babst, T. Kaspar, G. Acuña, F. Arigoni, and H. Hennecke. 1993. One member of a *groESL*-like chaperonin multigene family in *Bradyrhizobium japonicum* is co-regulated with symbiotic nitrogen fixation genes. *EMBO J.* 12:2901–2912.
- Frees, D., and H. Ingmer. 1999. ClpP participates in the degradation of misfolded protein in *Lactococcus lactis*. *Mol. Microbiol.* 1:79–87.
- Gaillot, G., E. Pellegrini, S. Bregenholt, S. Nair, and P. Berche. 2000. The ClpP serine-protease is essential for the intracellular parasitism of the human pathogen *Listeria monocytogenes*. *Mol. Microbiol.* 35:1286–1294.
- Gerth, U., E. Krüger, I. Derré, T. Msadek, and M. Hecker. 1998. Stress induction of the *Bacillus subtilis* *clpP* gene encoding a homologue of the proteolytic component of the Clp protease and the involvement of ClpP and ClpX in stress tolerance. *Mol. Microbiol.* 28:787–802.
- Gerth, U., A. Wipat, C. R. Harwood, N. Carter, P. T. Emmerson, and M. Hecker. 1996. Sequence and transcriptional analysis of *clpX*, a class-III heat-shock gene of *Bacillus subtilis*. *Gene* 181:77–83.
- Gething, M.-J., and J. Sambrook. 1992. Protein folding in the cell. *Nature* 355:33–45.
- Gottesman, S., S. Wickner, and M. R. Maurizi. 1997. Protein quality control: triage by chaperones and proteases. *Genes Dev.* 11:815–823.
- Grandvalet, C., V. de Crécy-Lagard, and P. Mazodier. 1999. The ClpB ATPase of *Streptomyces albus* G belongs to the HspR heat shock regulon. *Mol. Microbiol.* 31:521–532.
- Hamoen, L. W., A. F. Van Werkhoven, J. J. Bijlsma, D. Dubnau, and G. Venema. 1998. The competence transcription factor of *Bacillus subtilis* recognizes short A/T-rich sequences arranged in a unique, flexible pattern along the DNA helix. *Genes Dev.* 12:1539–1550.
- Hecker, M., W. Schumann, and U. Völker. 1996. Heat-shock and general stress response in *Bacillus subtilis*. *Mol. Microbiol.* 19:417–428.
- Herbort, M., U. Schön, K. Angermann, J. Lang, and W. Schumann. 1996. Cloning and sequencing of the *dnaK* operon of *Bacillus stearothermophilus*. *Gene* 170:81–84.
- Homuth, G., S. Masuda, A. Mogk, Y. Kobayashi, and W. Schumann. 1997. The *dnaK* operon of *Bacillus subtilis* is heptacistronic. *J. Bacteriol.* 179:1153–1164.
- Homuth, G., A. Mogk, and W. Schumann. 1999. Post-transcriptional regulation of the *Bacillus subtilis* *dnaK* operon. *Mol. Microbiol.* 32:1183–1197.
- Ingmer, H., F. K. Vogensen, K. Hammer, and M. Kilstrup. 1999. Disruption and analysis of the *clpB*, *clpC*, and *clpE* genes in *Lactococcus lactis*: ClpE, a new Clp family in gram-positive bacteria. *J. Bacteriol.* 181:2075–2083.
- Kobayashi, K., H. Wada, K. Asai, S. Moriya, and N. Ogasawara. 1999. Characterization of *ywlE* and *yfkJ* genes encoding low molecular weight protein-tyrosine phosphatases in *Bacillus subtilis*. Presented at 10th International Conference on Bacilli, Baveno, Italy.
- Krüger, E. 1998. Unpublished data.
- Krüger, E., and M. Hecker. 1998. The first gene of the *Bacillus subtilis* *clpC* operon, *ctsR*, encodes a negative regulator of its own operon and other class III heat shock genes. *J. Bacteriol.* 180:6681–6688.
- Krüger, E., T. Msadek, and M. Hecker. 1996. Alternate promoters direct stress-induced transcription of the *Bacillus subtilis* *clpC* operon. *Mol. Microbiol.* 20:713–724.

71. Schulz, A., S. Schwab, S. Versteeg, and W. Schumann. 1997. The *htpG* gene of *Bacillus subtilis* belongs to class III heat shock genes and is under negative control. *J. Bacteriol.* 10:3103–3109.
72. Schulz, A., B. Tzschaschel, and W. Schumann. 1995. Isolation and analysis of mutants of the *dnaK* operon of *Bacillus subtilis*. *Mol. Microbiol.* 15:421–429.
73. Schumann, W. 1999. FtsH—a single-chain charonin? *FEMS Microbiol. Rev.* 23:1–11.
74. Schweder, T., K. H. Lee, O. Lomovskaya, and A. Martin. 1996. Regulation of *Escherichia coli* starvation sigma factor (σ^s) by ClpXP protease. *J. Bacteriol.* 178:470–476.
75. Servant, P., C. Grandvalet, and P. Mazodier. 2000. The RheA repressor is the thermosensor of the HSP18 heat shock response in *Streptomyces albus*. *Proc. Natl. Acad. Sci. USA* 97:3538–3543.
76. Servant, P., G. Rapoport, and P. Mazodier. 1999. RheA, the repressor of *hsp18* in *Streptomyces albus* G. *Microbiology* 145:2385–2391.
77. Tozawa, Y., H. Yoshikawa, F. Kawamura, M. Itaya, and H. Takahashi. 1992. Isolation and characterization of the *groES* and *groEL* gene of *Bacillus subtilis* Marburg. *Biosci. Biotechnol. Biochem.* 56:1995–2002.
78. Turgay, K., J. Hahn, J. Burghoorn, and D. Dubnau. 1998. Competence in *Bacillus subtilis* is controlled by regulated proteolysis of a transcription factor. *EMBO J.* 17:6730–6738.
79. Turgay, K., L. W. Hamoen, G. Venema, and D. Dubnau. 1997. Biochemical characterization of a molecular switch involving the heat shock protein ClpC, which controls the activity of ComK, the competence transcription factor of *Bacillus subtilis*. *Genes Dev.* 11:119–128.
80. Van Asseldonk, M., A. Simons, H. Visser, W. M. De Vos, and G. Simons. 1993. Cloning, nucleotide sequence, and regulatory analysis of the *Lactococcus lactis* *dnaJ* gene. *J. Bacteriol.* 175:1637–1644.
81. Vanet, A., J. A. Plumbridge, and J.-H. Alix. 1993. Co-transcription of two genes necessary for ribosomal protein L11 methylation (*prmA*) and pantothenate transport (*panF*) in *Escherichia coli* K-12. *J. Bacteriol.* 175:7178–7188.
82. Van Sinderen, D., A. Luttinger, L. Kong, D. Dubnau, G. Venema, and L. W. Hamoen. 1995. *comK* encodes the competence transcription factor, the key regulatory protein for competence development in *Bacillus subtilis*. *Mol. Microbiol.* 15:455–462.
83. Varmanen, P., H. Ingmer, and F. K. Vogensen. 2000. *ctsR* of *Lactococcus lactis* encodes a negative regulator of *clp* gene expression. *Microbiology* 146:1447–1455.
84. Versteeg, S., and W. Schumann. Unpublished data.
85. Versteeg, S., A. Mogk, and W. Schumann. 1999. The *Bacillus subtilis* *htpG* gene is not involved in thermal stress management. *Mol. Gen. Genet.* 261:582–588.
86. Wehrli, W., M. Niederweis, and W. Schumann. 2000. The FtsH protein accumulates at the septum of *Bacillus subtilis* during cell division and sporulation. *J. Bacteriol.* 182:3870–3873.
87. Wetzstein, M., U. Völker, J. Dedio, S. Löbau, U. Zuber, M. Schiesswohl, C. Herget, M. Hecker, and W. Schumann. 1992. Cloning, sequencing, and molecular analysis of the *dnaK* locus from *Bacillus subtilis*. *J. Bacteriol.* 174:3300–3310.
88. Wiegert, T., and S. Zellmeier. Personal communication.
89. Wiegert, T., and W. Schumann. Unpublished work.
90. Yuan, G., and S.-L. Wong. 1995. Isolation and characterization of *Bacillus subtilis* regulatory mutants: evidence for *orf39* in the *dnaK* operon as a repressor gene in regulating the expression of both *groE* and *dnaK*. *J. Bacteriol.* 177:6462–6468.
91. Yuan, G., and S.-L. Wong. 1995. Regulation of *groE* expression in *Bacillus subtilis*: the involvement of the σ^A -like promoter and the roles of the inverted repeat sequence (CIRCE). *J. Bacteriol.* 177:5427–5433.
92. Yura, T., M. Kanemori, and M. Morita. 2000. The heat shock response: regulation and function, p. 3–18. In G. Storz and R. Hengge-Aronis (ed.), *Bacterial Stress Response*. American Society for Microbiology, Washington, D.C.
93. Zuber, U., K. Drzewiecki, and M. Hecker. Unpublished data.
94. Zuber, U., and W. Schumann. 1994. CIRCE, a novel heat shock element involved in regulation of heat shock operon *dnaK* of *Bacillus subtilis*. *J. Bacteriol.* 176:1359–1363.